

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

ANTIBODY, ANTIBODY FORMING CELL AND PRODUCTION OF ANTIBODY

Patent Number: JP2069196
Publication date: 1990-03-08
Inventor(s): MIYAZAKI KIMIMASA; others: 02
Applicant(s): MATSUSHITA ELECTRIC IND CO LTD
Requested Patent: ☐ JP2069196
Application Number: JP19880220932 19880902
Priority Number(s):
IPC Classification: C12P21/08; A61K39/395; C12N5/16; C12N15/06; G01N33/531; G01N33/577
EC Classification:
Equivalents: JP2600325B2

Abstract

PURPOSE: To obtain high performance measurable in high sensitivity by making pseudo-antigen in which alkyl group is substituted by H-atom to immunogen by chemically bonding with protein.
CONSTITUTION: N-Alkyl group in antigen of low-molecular weight compound having ≤ 2000 molecular weight such as methamphetamine (MA) expressed by formula I is substituted with H-atom to obtain amphetamine (AP) expressed by formula II and aminobutyl group is introduced into said amphetamine as spacer to obtain antigen derivative (ABAP) expressed by formula III. Next, said ABAP is bonded with hemocyanin (KLH) derived from false limpet as carrier protein to obtain immunogen AP-KLH. Then, said immunogen is injected into animal such as system A/J of mouse and antibody such as anti-MA antibody is produced in blood serum. Cell of sensitized animal such as said immuno-injected mouse is fused with myeloma cell to obtain antibody forming cell and said antibody forming cell is cultured in medium, as necessary, then resultant cultured supernatant is purified to collect monoclonal antibody.